



XI Simpósio Brasileiro de Farmacognosia
 XVI Simposio Latinoamericano de Farmacobotanica
 Curitiba, Brasil - 9-11 de agosto de 2017

Biotransformation and evaluation of the leishmanicidal activity of quinolinic alkaloids and derivatives

**Keylla Lençone Mischiatti,¹ Brás Heleno de Oliveira,¹ Almeriane Weffort-Santos²,
 Lucélia Donatti³**

¹Departamento de Química, Universidade Federal do Paraná, UFPR, Curitiba/PR, Brazil

²Departamento de Análises Clínicas, Universidade Federal do Paraná, UFPR, Curitiba/PR, Brazil

³Departamento de Biologia Celular, Universidade Federal do Paraná, UFPR, Curitiba/PR, Brazil

Introduction: In recent years new compounds with anti-leishmanial activity derived from quinolinic alkaloids have been developed. **Objectives:** The aim of this work was to produce phenylquinoline alkaloid (PQ) derivatives by biotransformation using *Aspergillus fumigatus* and *A. flavus*, and compare their leishmanicidal potential with the original molecule. **Material & Methods:** The microorganisms' cultures were shaken (150 rpm) for 7 days at 28 °C, in medium (200 ml) enriched with glucose, malt and yeast extracts added of PQ (100 mg). The products were purified by chromatographic techniques and identified by MASS and NMR spectroscopy. **In vitro** leishmanicidal activity against *L. amazonensis*, *L. infantum*, and *L. braziliensis* promastigote forms of PQ and derivatives were performed using the MTT assay, and photodocumented by transmission electronic microscopy (TEM). The effects were compared with standard drugs. **Results:** Two derivatives were obtained: DFQ1, from *A. fumigatus*; 8% yield, with two hydroxyl groups at C5 and C6 positions, originating a substance not reported to date; and DFQ2, from *A. flavus*; 5% yield, with a hydroxylation at the C4' position. Table 1 shows the leishmanicidal activity, with IC₅₀ results according to the incubation time. Interestingly, the extra hydroxyl groups of DFQ1 enhanced the activity of the original molecule. TEM of the parasites showed mitochondrial swelling, increase of lipid inclusions, chromatin condensation, and changes in the plasma membrane. **Conclusion:** The results herein presented showed that PQ could be biotransformed by *A. fumigatus* producing DFQ1, a new unreported substance. *A. flavus* also produced DFQ2, and both quinolinic derivatives showed higher leishmanicidal potential than the parental molecule disrupting directly the cytoplasmatic, nuclear, and membrane components of the parasites.




Figure 1. NMR¹H [400 MHz, (CD₃)₂CO] of DFQ1 and DFQ2.

Table 1. Leishmanicidal activity of phenylquinoline (PQ) and derivative alkaloids.

SUBSTANCE	Incubation time (h)	<i>L. amazonensis</i>	<i>L. braziliensis</i>	<i>L. infantum</i>
		IC ₅₀ (μM)	IC ₅₀ (μM)	IC ₅₀ (μM)
PQ	24	>500	>500	>500
	48	>500	>500	>500
	72	>500	>500	>500
DFQ1	24	5,41±0,20	98,25±0,22	69,54±0,19
	48	0,44±0,18	63,01±0,28	7,37±0,23
	72	0,24±0,14	10,32±0,26	4,27±0,32
DFQ2	24	298,64±0,35	385,65±0,71	290,67±0,34
	48	203,65±0,70	289,32±0,57	230,45±0,45
	72	165,45±0,35	202,65±0,35	178,42±0,71
Anfotericin B	24	285,60±0,35	345,65±0,28	315,64±0,85
	48	125,50±0,45	142,70±0,47	67,46±0,38
	72	79,40±0,48	122,80±0,60	41,48±0,37
Glucantime	24	12,50±0,41	20,06±0,50	19,88±0,48
	48	9,60±0,43	16,78±0,51	8,72±0,54
	72	7,84±0,42	15,39±0,53	6,95±0,57

Keywords: biotransformation, quinolinic alkaloids, leishmanicidal activity.